

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

QUARTERLY (First)

REPORT ON

(15) CONTRACT ~~NO~~ DA 92/5574FEC-34627

INCLUSIVE DATES 1 February TO 30 April, 1961

Page # M23-59-23
64-13

①

DI in 1000 (1, 2, 3, 4, 5, 6, 7, 8, 9, 10)

SUBJECT OF INVESTIGATION

CORRELATION BETWEEN SUSCEPTIBILITY TO
ORAL INFECTION AND INTESTINAL BACTERIAL
FLORA IN THE INBRED MOUSE STRAINS RAISED
BY SELECTIVE BROTHER-SISTER MATINGS IN
OUR LABORATORY

RESPONSIBLE INVESTIGATOR

Shogo Sasaki, M.D.

Assistant Professor of Bacteriology
School of Medicine, Keio University
Shinjuku-ku, Tokyo, Japan

U.S. Army Research & Development Group(9852)(Far East)

Office of the Chief of Research and Development

United States Army

APO 343

AD No. ~~1~~
DBC FILE COPY

438691

438691

QUARTERLY (First)

92-557-FEC-34627

1 February

30 April

Correlation between susceptibility to
oral infection and intestinal bacterial flora
in the inbred mouse strains raised by brother-
sister matings in our laboratory.

Shogo Sasaki, M. D.

Assistant Professor of Bacteriology
School of Medicine, Keio University
Shinjuku-ku, Tokyo, Japan

Correlation between susceptibility to oral infection
and intestinal bacterial flora in the inbred mouse
strains raised by brother-sister matings in our laboratory.

Shogo Sasaki, M. D.

Assistant Professor of Bacteriology
School of Medicine, Keio University
Shinjuku-ku, Tokyo, Japan

SCOPE OF SERVICES

1) Mass breed two inbred mouse strains.

We have managed to raise three inbred mouse strains by selective brother-sister matings started from one pair of "ddN" strain mice. These strains, which we named dd/Kf, dd/Ks, and dd/Kr, have been selected for their susceptibility to infection with Salmonella enteritidis, and they are different from each other not only in susceptibility to the infection, but also in immunizability against it.

Two of these strains, dd/Ks and dd/Kr, was selected for this research. The dd/Ks strain mice have a higher susceptibility to the salmonella infection than dd/Kr strain mice.

We started to produce monthly 150 disease-free mice of the two strains. For this purpose 60 females and 60 male mice was provided as parents based on the assumption that the size of a litter will be 5.

2) Study the relation between normal intestinal flora to the infection.

We have been studying oral infection in experimental animals and found that multiplication of infected microorganisms in the intestinal tract had a close relation to the normal

intestinal bacterial flora of the host. The results as follow: adult mice had more stable intestinal flora than baby and orally administered bacteria were eliminated from the intestines within a short period and their own normal flora did not change at all. After the derangement or elimination of normal intestinal flora by X-irradiation, starvation, administration of cortisone , OCl₄ , or some antibiotics newly administered bacteria resided in the intestines for a long time.

Based on these lines of basic study we intend to study systematically the intestinal flora of these inbred mouse strains to clarify more precisely the significans of each member of intestinal flora to oral infection with Salmonella or Shigella.

a. Significance of Enterococci as normal intestinal flora to the infection. As the representative of gram positive cocci in normal intestinal flora, we selected Enterococcus and gave them orally after pretreatment with antibiotics to clarify the significans of this bacteria to the following oral infection with Salmonella or Shigella.

b. Significance of gram negative bacilli as normal intestinal flora to the infection. For the same purpose as mentioned above, one strain of Escherichia coli was selected as the representative of gram negative bacilli in normal intestinal

flora.

We are expecting that the correlation between orally infected Salmonella or Shigella, and Enterococci or gram negative bacilli which are both the members of normal intestinal flora, will be clarified by these experiments.

3) Examine genetical characteristics of each inbred mouse strain by reciprocal matings and cellular and humoral factors making up the host's defense to infection of the mice.

The purpose of this experiment is to test the homogeneity of genetic constitution of these strains.

As one of the tests to prove the homogeneity cell-transfer will be carried out.